

Product Number: 62131
ATS Number: 49196
Pack Sizes: 1L-1000L

62131_49196_500mL-1000L_primary_MPL_V2
Label Size: 165mm x 175mm
PDF Scale: 100% A4 / 100% A3

ACS SWAN EXPRESS PRINT

Filename: 43695_4F_Propyzamide500_front
Modified: July 9, 2010 1:31 PM

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



PROPYZAMIDE 500SC Herbicide

ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE

GROUP D HERBICIDE

For selective control of certain grasses and broad leaf weeds in lettuce, sports turf, home lawns and legume seed crops and pastures, as per the Directions for Use Table

IMPORTANT: READ ALL INSTRUCTIONS BEFORE USING THIS PRODUCT.

Contents: 20 Litres



4FARMERS PTY. LTD.

A.C.N. 067 443 485

1/70 McDOWELL ST, WELSHPOOL,
WESTERN AUSTRALIA 6106

PH: (08) 9356 3445 FAX: (08) 9356 3447

Product Number: 62131
ATS Number: 49196
Pack Sizes: 1L-1000L

62131_49196_500mL-1000L_immediate_MPL_V2
Label Size: 250mm x 175mm
PDF Scale: 99% A4 / 100% A3

ACS SWAN EXPRESS PRINT

Filename: 43695_4F_Propyzamide500_Back2

Modified: July 9, 2010 1:25 PM

DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate L/ha	Critical Comments
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All states	4.5 in 330-550 L	Apply a pre-emergent immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
Transplanted				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
		Vic only	2.3 in 300-550 L	Apply on light sandy soils, low in organic matter.
Turf & Lawns Common Couch	Winter Grass	All states	1.2 (12 mL/100m ²)	Rate/Bowling Green 37 x 37 m 170 mL Apply during the autumn and winter, if established turf is infested with seedling winter grass.
		Old only	0.6 (6 mL/100 m ²)	85 mL Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Old blue couch after 4 weeks.
Buffalo Grass		WA only	1.2 (12 mL/100 m ²)	170 mL
Lucerne, Clover, Medics etc. Grown for Forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed	NSW, ACT, SA, WA, Tas only	1-1.5 1.5-2	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed ESTABLISHED CROPS: Apply anytime before end of July. WEEDS: Apply when weeds are at post or Pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of Propyzamide 500SC to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

Resistant Weeds Warning

Propyzamide 500SC Herbicide is a member of the Benzamide group **GROUP D HERBICIDE** of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management this is a group D herbicide. Some naturally occurring weed biotypes resistant to Propyzamide 500SC Herbicide and other group D herbicides may exist through normal genetic variability in any weed population. The resistant individual can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Propyzamide 500SC Herbicide or other group D Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result in the failure of Propyzamide 500SC Herbicide to control the resistant weeds. Advice as to strategies and alternative and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Sipcarn representative.

PROTECTION OF WILDLIFE,

FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, and waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS,

NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plant/crops, cropping lands or pastures.

STORAGE AND DISPOSAL

(500 ml to 20 L)

Store in a closed original container in a cool, dry, well-ventilated area out of direct sunlight. Do not contaminate seed, feed or foodstuff. Do not re-use container for any purpose. Triple or preferably pressure rinse containers before disposal. Add rinsings or spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler of designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers to below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Do not store for prolonged periods in direct sunlight.

Refillable containers

(1000 L only)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

4Farmers Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use or abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on 4Farmers judgment in purchasing or using the product and every person dealing with this product does so at their own risk.

**ONLY IN EMERGENCY CALL
DIAL 000
FOR POLICE OR THE FIRE BRIGADE
ALL HOURS**

Batch Number:

Date of Manufacture:



APVMA Approval No: 62131/1L-1000L/0210